This paper can be downloaded at the web site of the Japanese Journal of Applied Physics, http://www.ipap.jp/jjap/index.htm. Until January 2004, anyone could register and download papers there at no cost. The journal is now charging for reprints. Here is the title, abstract, keywords and Fig. 1 from this paper:

Jpn. J. Appl. Phys. Vol. 38(1999) L774-L776

Part 2, No. 7A, 1 July 1999 DOI: 10.1143/JJAP.38.L774

Observation of Anomalous Heat Release and Helium-4 Production from Highly Deuterated Palladium Fine Particles

Yoshiaki Arata and Yue-Chang Zhang

Cooperation Research Center for Advanced Science and Technology, Osaka University, 11-1 Mihogaoka, Ibaraki, Osaka 567-0047, Japan

(Received April 22, 1999; revised manuscript received May 17, 1999; accepted for publication May 19, 1999)

Abstract:

Observations were made of the anomalous production of 4 He atoms as well as the anomalous heat release when Pd fine particles are highly deuterated inside an enclosed Pd metal vessel used as a cathode in electrolysis of D_2O . A mass analysis of the remnant Pd powders after the 2000-hr heat production revealed substantial production of 4 He atoms.

Keywords: double-structure cathode, deuterated Pd, Pd black, electrolysis, excess energy, ⁴He production




Fig. 1. (Left) Schematic view of the electrolysis apparatus with a DS-cathode. The DS-cathode is surrounded by a cylindrical Pt anode. The electrolyte (water+0.1 M LiOH) is continuously cooled by chilled water flowing through a cooling coil. (Right) Structure of the DS-cathode used for the present experiments. Fine particles of Pd (Pd black) of about 20 nm diameter are enclosed in a vacuum in a Pd metal vessel of 3 mm wall thickness, 14 mm outer diameter and 50–70 mm length.